



## New Approaches for Prevention from Musculoskeletal Pains

Ali Ghanjal

Health Management Research Centre, Department of Physical Medicine and Rehabilitation, Baqiyatallah University of Medical Sciences, Tehran, Iran.

Musculoskeletal pains are a common symptom in many people (Malmberg-Ceder et al., 2017). In Europe, musculoskeletal pains account for approximately 40% of all occupational diseases and are considered as a growing problem (European Agency for Safety and Health at Work, 2017). The presence of musculoskeletal pains has been associated with reduced quality of life for individuals, decreased productivity, increased sickness, absence from the workplace, and economic consequences for the society (European Agency for Safety and Health at Work, 2017; McDonald, Dibonaventura, & Ullman, 2011; Cancelliere et al., 2011).

Musculoskeletal pain is usually caused by: Performing repetitive activities in inappropriate body conditions, Keeping body in a steady state for a long time, not paying attention to ergonomic issues in daily activities and during work, not performing simple and periodic exercises during daily activities, having poor physical fitness for exercise, having weak muscles, shortening the length of the tendons and ligaments and fascia, doing improperly stretching exercises, Removing and moving heavy loads (especially high altitude), not paying attention to psychological issues and (Dalager, Justesen, & Sjøgaard, 2017)

Therefore, in order to prevent from the occurrence of these pains, special attention should be paid to the following approaches:

Changing the status of the body during activities; performing simple and periodic exercises; paying attention to stretching exercises, to ergonomics in life and work, to doing daily activities in a proper physical condition, to the issue of physical fitness in carrying out activities; reducing repetitive tasks; Intervention in the prevention from musculoskeletal pains; early Identification (Dalager, Justesen, & Sjøgaard, 2017; Linton, 2002; Falla et al., 2017); and the use

of virtual reality in the treatment of musculoskeletal pains (manipulating sensory signs to improve motor function during walking) (Powell & Simmonds, 2014).

### References

- Malmberg-Ceder, K., Haanpää, M., Korhonen, P. E., Kautiainen, H., Soinila, S. (2017). Relationship of musculoskeletal pain and well-being at work-Does pain matter? *Scandinavian Journal of Pain*, 15, pp. 38-43.
- European Agency for Safety and Health at Work. OSH in figures: Work-related musculoskeletal disorders in the EU: Facts and figures. 2017, p. 1-184.
- McDonald, M., Dibonaventura, M. D., & Ullman, S. (2011). Musculoskeletal pain in the workforce: The effects of back, arthritis, and fibromyalgia pain on quality of life and work productivity. *Journal of Occupational and Environmental Medicine*, 53 (7), pp. 765-770.
- Cancelliere, C., Cassidy, J. D., Ammendolia, C., & Côté, P. (2011). Are workplace health promotion programs effective at improving presenteeism in workers? A systematic review and best evidence synthesis of the literature. *BMC Public Health*, 11 (1), p. 395 .
- Dalager, T., Justesen, J. B., & Sjøgaard, G. (2017). Intelligent physical exercise training in a workplace setting improves muscle strength and musculoskeletal pain: A randomized controlled trial. *BioMed Research International*, 2017(2017), pp. 1-9.
- Linton, S. J. (2002). Early identification and intervention in the prevention of musculoskeletal pain. *American Journal of Industrial Medicine*, 41 (5). pp. 433-442.
- Falla, D. I., Cescon, C., Lindstroem, R., Barbero, M. (2017). Muscle pain induces a shift of the spatial distribution of upper trapezius muscle activity during a repetitive task: A mechanism for perpetuation of pain with repetitive activity? *The Clinical Journal of Pain*.
- Powell, W., Simmonds, M. J. (2014). Virtual reality and musculoskeletal pain: Manipulating sensory cues to improve motor performance during walking. *Cyberpsychology Behavior and Social Networking*, 17 (6): 390-6.