

## Fitness Enhancement by Crosses between Two Populations of *Trissolcus vassilievi* (Hymenoptera: Scelionidae)

Shahzad Iranipour<sup>1</sup>, Parisa Benamolaei<sup>2\*</sup>, and Shahriar Asgari<sup>3</sup>

### ABSTRACT

*Trissolcus vassilievi* (Mayr) (Hymenoptera: Scelionidae) is one of the most important egg parasitoids of the Common Sunn Pest (CSP), *Eurygaster integriceps* Puton (Hemiptera: Scutelleridae) in Iran. In this study, the fitness of two populations of *T. vassilievi* was studied on two populations of hosts in terms of life history parameters. Two populations of both *T. vassilievi* and the host, CSP were selected: (1) Tabriz (as a temperate area), and (2) Varamin (as a subtropical area), for CSP. Moreover, regarding that outcrossing between populations can produce progeny with superior characteristics, the progeny of reciprocal crosses between original populations also were examined on a single host. The crosses between the two populations caused 13.9-18.5% higher net fecundity than maternal populations, which suggests fecundity to be a function of maternal phenotype. The intrinsic rate of increase showed minor differences among treatments, which varied between  $0.291 \pm 0.003$  to  $0.305 \pm 0.003$ . The partial advantage of Varamin wasps over Tabriz, and the crosses over the original populations was obvious. Such differences may be used to obtain more efficient parasitoids in augmentation programs.

**Keywords:** Intrinsic rate of increase, Life expectancy, Net reproduction rate, Parasitoid wasp, Reciprocal crosses.

### INTRODUCTION

The biological fitness of a living organism is the relative ability of an organism to survive and pass on its genes to the next generation (Krebs and Davies, 1993). This is possible through birth (transferring more proportion of genes to the next generation) in the shortest possible time (high speed of gene transcription) and the ability of progeny to survive (persistence of the genes). Therefore, the developmental rate, the mortality rate at all stages of life, and fecundity are relevant features of fitness and the ability of a living organism to compete with other species. These characteristics can

be examined as life tables and stable population growth models (Lotka, 1907a, b; Portilla *et al.*, 2014). So far, several studies have investigated the stable population growth parameters in egg parasitoids of Common Sunn Pest (CSP), *Eurygaster integriceps* Puton (Hem., Scutelleridae), and related stinky bugs. These are dominantly on *Trissolcus* spp. and *Telenomus* spp. (Hymenoptera: Scelionidae) (Asgari and Kharrazi Pakdel, 1998; Laumann *et al.*, 2008; Amir-Maafi and Parker, 2011; Nozad-Bonab *et al.*, 2014; Bazavar *et al.*, 2015; Benamolaei *et al.*, 2015a, b; Abdi *et al.*, 2017; Teimouri *et al.*, 2019), and *Ooencyrtus* spp. (Hymenoptera: Encyrtidae)

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, University of Tabriz, Tabriz, Islamic Republic of Iran.

<sup>2</sup> Department of Animal Biology, Faculty of Natural Science, University of Tabriz, Tabriz, Islamic Republic of Iran.

<sup>3</sup> Tehran Agricultural and Natural Resources Research and Education Center, AREEO, Varamin, Islamic Republic of Iran.

\*Corresponding author; e-mail: p.benamolaei@tabrizu.ac.ir



(Ahmadpour et al., 2013; Mele et al., 2024).

There is little information about intra- and inter-specific variation of survival value and reproductive potential of egg parasitoids of CSP, and it is limited to few studies. In the 1970s, several species of CSP's egg-parasitoids, collected from different regions of the world, were transferred to the former Soviet Union and compared with native species, especially *Trissolcus grandis* (Thomson), in laboratory and field conditions. The overall result of this study showed that native species had superiority and it was due to the adaptation of native species to climatic and seasonal conditions (Nouri et al., 2011). Awan et al. (1990) compared three geographical populations of *T. basalis* (Wollaston) collected from France, Italy, and Spain regarding biological and behavioral characteristics. The emergence rate of adult wasps from *Nezara viridula* (L.) eggs was significantly higher in the French population than in the Italian and Spanish ones. The development of immature stages of the Italian population was significantly longer than the other two populations. Taghadosi et al. (1993) and Nozad Bonab et al. (2014) observed differences among populations of *T. grandis* of Tehran-Alborz-Qazvin and East Azarbaijan provinces, respectively. A comparison of biostatistics of Scelionidae by Amir-Maafi (2010) during 2004-2006 in different provinces of Iran revealed significant differences between species and populations of the wasps. Fecundity, oviposition period, and gross and net reproductive rates differed between species or populations. The net fecundity of *T. vassilievi* (Mayr) from Lorestan and Tehran was 240.8 and 227.5 eggs, respectively, which was 2.5 times the other populations. The highest intrinsic rate of increase of *T. grandis*, *T. semistriatus* (Nees), and *T. vassilievi* was recorded for Golestan, West Azarbaijan, and Tehran provinces, respectively.

It can be seen from the above reports that inter-specific and intra-specific differences in the parasitism rate and population growth

rate of the parasitoid wasps are sometimes very considerable. Therefore, in this study, we attempted to study the differences between two populations of *T. vassilievi* with temperate (Tabriz) and subtropical (Varamin) origins simultaneously. On the other hand, since the parasitoid spends all immature stages within the host body, it can be affected by the host's quality, so the host population was also included as a second variable. Finally, the hypothesis was tested to find if crosses between populations could enhance the fitness of populations. Therefore, crosses as Tabriz-female×Varamin-male (T×V) and Varamin-female×Tabriz-male (V×T) were conducted to evaluate the possibility of obtaining populations with superior or intermediate characteristics.

## MATERIALS AND METHODS

### Cultures of *Eurygaster integriceps* Puton

Adult bugs were collected on several occasions at the end of the winter from mountains around Tabriz and Varamin before leaving the resting sites. Collecting of specimens were continued in wheat fields during post diapause phase. The collected insects were transferred to a greenhouse unit of the Department of Plant Protection, Faculty of Agriculture, University of Tabriz. Transparent rectangular cubic plastic containers (20×30×9 cm) equipped with a mesh cap for ventilation were used for the rearing of both populations. Dry wheat grain was used as a foodstuff and soaked cotton balls as a water source. The paper strips folded fan-like to serve as an oviposition substrate. These insects were exposed to 25±2°C, 40±10% RH, and 16:8 h L:D photoperiod (Iranipour et al. 2015).

### Cultures of *Trissolcus vassilievi* (Mayr)

In this study, two original populations of *T. vassilievi*, namely, one from Tabriz (1360

AMSL, 46°E, 38°N) and the other from Varamin (918 AMSL, 51°E, 35°N) were examined. To collect egg parasitoids, host egg traps (yellow cardboards, 5×15 cm, folded twice to construct a  $\Delta$ -shaped structure) were used (Safavi, 1973). The traps were tied to wheat ears and removed after one week. Then, the parasitized eggs were transferred to glass vials (1.5×10 cm) and kept in a growth chamber (Iran Khodsaz Co., IKH.RH model) under constant conditions (26±1°C, 50±5% RH and 16:8 h L:D photoperiod). The emerged wasps were identified by the identification key of Kozlov and Kononova (1983). After rearing for one generation, males and females of the second generation from the same population were randomly coupled and each pair was transferred to a similar vial supplied by the host eggs of either population. Small drops of honey were used to feed the wasps.

### Fecundity-Life Table Studies

Ten clutches of 24-hour-old CSP eggs (14 eggs per clutch) from each population were exposed to 48-hour-old mated *T. vassilievi* females of the second-generation of each population. After 24 hours, females were removed and the host eggs were kept as a life table cohort in the so-called growth chambers to determine their fate. The experiment was conducted as a factorial experiment in a completely randomized design framework, with two factors including wasp and host populations, respectively; each one in two levels that were represented as TT, TV, VT, and VV (original populations); where, the first letter delineates the parasitoid origin, and the second one the host origin (T= Tabriz, V= Varamin). Twenty pairs of third-generation wasps from each population were coupled randomly, and five clutches of 24-hour-old CSP eggs were offered daily until death. The fate of host eggs was followed by daily checks, and the date of emergence was recorded separately for males and females.

In the next step, the progeny of the third generation of the two populations was reciprocally crossed to the other population.

Thus, two kinds of the cross were present: Tabriz females×Varamin males (T×V) and Varamin female×Tabriz males (V×T). Considering the non-significant effect of the host, the outcrossed wasps were studied only on CSP eggs of Varamin in the same manner described for original populations.

### Measures of Stable Population Growth Parameters

The method described by Carey (1993) was used to calculate life table parameters and entropy (Equation 1).

$$H = \frac{\sum_{x=0}^{\omega} e_x d_x}{e_0} \quad (1)$$

$e_0$ ,  $e_x$ , and  $d_x$  represent life expectancy at age 0, and life expectancy and mortality distribution at age  $x$ , respectively, summed over ages 0 to  $\omega$  (the last death event), and  $H$  denotes entropy.

The entropy values less than, equal to, and greater than 0.5 represent the survivorship curves of type I (convex), type II (straight line), and type III (concave), respectively. Stable population growth parameters, including Gross Reproduction Rate (GRR), net Reproduction rate ( $R_0$ ), mean generation Time (T), Doubling Time (DT), intrinsic rate of increase ( $r_m$ ), finite rate of increase ( $\lambda$ ), intrinsic birth rate (b), and intrinsic death rate (d) were estimated. The  $r_m$ -value was calculated by solving the Lotka equation using iterative calculations of the Newton-Raphson method as follows:

$$\sum_{x=0}^{\omega} e^{-r_m x} l_x m_x = 1 \quad (2)$$

$l_x$  and  $m_x$  represents survival rate and fecundity at age  $x$  respectively,  $r_m$  denotes intrinsic rate of increase, and  $\sum$ ,  $\omega$ , and  $e$  are symbols of summation, last event of cohort and natural logarithm base respectively.

To determine the standard error of the above statistics, we used bootstrap methods in 1000 replicates (Meyer *et al.*, 1986). Estimation of the parameters was carried out by a program in Excel (Iranipour, 2018).



## Statistical Analyses

Statistical analysis was performed using SPSS software. Since the host-effect was non-significant in all parameters, this factor was excluded from analysis and comparison between main populations and crossed populations was done by One-Way ANOVA. The means were compared by Tukey's test at 0.05 significance level. Bootstrap estimates of  $r_m$  and the other stable population growth parameters were also compared among the main populations and crosses. Pairwise comparisons between those treatments were done by the random pairing of estimates, and the differences of randomly paired values were ranked from the smallest to the largest value. Then 25 boundary values from both ends were excluded and 950 median values were considered as 95% Confidence Interval (CI). If 95% CI included zero, no significant difference between the two treatments was interpreted.

## RESULTS

### Life History Parameters of *T. vassilievi*

In life table studies, two populations of *T. vassilievi* were compared on two host populations (Table 1), and due to the insignificant effect of the host and their interactions, the host effect was excluded from the analysis. The wasps from the crosses on the Varamin host were compared with their parental populations only on the Varamin host.

Analysis of variance showed that there was a significant difference in the longevity of both females and males when crosses were included

as well (Tabriz population, Varamin population, T×V, and V×T crosses;  $F_{3,76} = 3.62$ ,  $P = 0.017$  for female longevity, and  $F_{3,76} = 3.12$ ,  $P = 0.031$  for male longevity). The highest longevity of females and males was observed in Tabriz and Varamin wasps, respectively (Table 2).

Female wasps from the crosses showed intermediate longevity of the two populations, so, their differences were not significant with either population. However, the male progeny of the crosses had a shorter lifespan than their parents (Table 2). Females lack a pre-oviposition period. The mean oviposition period of *T. vassilievi* females was significantly higher in the Varamin population than in Tabriz and cross populations ( $F_{3,76} = 4.36$ ,  $P = 0.007$ ). The total fecundity of *T. vassilievi* females was significantly higher in cross populations than those of parental populations ( $F_{3,76} = 22.38$ ,  $P < 0.001$ ). Females of crosses laid 8–29% more eggs than both populations. The highest fecundity was observed in T×V. Almost in all cases, maximum oviposition occurred on the first day of life and declined to zero with a non-linear trend in less than three weeks (Figure 1). The highest fecundity was obtained in T×V, and V×T crosses, respectively, followed by the original populations (Table 2). The age-specific sex ratio (proportion of females) declined at senescence (Figure 2).

### Life Tables and Survivorship Curves of *T. vassilievi*

Age-specific mortality ( $q_x$ ) of *T. vassilievi* in all treatments increased by age. The results showed that mortality seldom occurs during and prior to oviposition. The survivorship

**Table 1.** Analysis of variance of life history components of the two populations of *T. vassilievi* on two populations of sunn pest eggs.<sup>a</sup>

Parameter	Wasp		Host		Wasp × Host	
	F	P	F	P	F	P
Female longevity	14.61	< 0.001	1.74	0.191	0.42	0.519
Male longevity	3.43	0.068	0.00	1.000	0.01	0.939
Oviposition period	29.41	< 0.001	0.31	0.580	0.17	0.678
Post-oviposition period	36.66	< 0.001	1.54	0.219	0.31	0.578
Total fecundity	24.16	< 0.001	1.80	0.183	1.58	0.212
Average daily fecundity	0.00	0.949	2.37	0.128	0.84	0.364
Sex ratio	0.04	0.847	0.01	0.932	0.22	0.641


<sup>a</sup> df for all treatments= 1,76.




**Table 2.** Reproductive parameters of two *T. vassilievi* populations on two CSP populations and crosses as T×V and V×T.<sup>a</sup>

Treatments	TT	VT	TV	VV	T×V	V×T
Female longevity (d)	31.80±1.19A	28.25±1.14B	31.05±1.15Aa	26.05±0.86Bb	29.90±1.50ab	28.10±1.23ab
Male longevity (d)	19.60±0.49A	20.75±0.55A	19.55±0.41Aab	20.80±0.95Aa	18.30±0.32b	18.75±0.32ab
Oviposition period (d)	18.50±0.41A	20.60±0.37B	18.45±0.39Ab	20.25±0.23Ba	18.40±0.48b	18.60±0.53b
Post oviposition period (d)	13.30±1.08A	7.65±1.15B	12.60±1.02Aa	5.80±0.84Bb	11.50±1.39a	9.50±1.01ab
Life time fecundity	216.90±5.36A	249.30±5.07B	230.55±6.02Ac	249.75±4.30Bb	280.40±2.77a	269.50±4.80a
Daily egg	11.80±0.34A	12.13±0.23A	12.63±0.49Ab	12.34±0.20Ab	15.46±0.47a	14.67±0.43a
Sex ratio	0.82±0.03A	0.82±0.02A	0.82±0.02Aa	0.82±0.02Aa	0.78±0.05a	0.86±0.01a

<sup>a</sup> Means bearing the same letter in a row are not significantly different (Tukey's HSD,  $\alpha=0.05$ ). Capital letters are for comparison between original populations and lower cases are for comparison between original populations and reciprocal crosses reared on the Varamin host.



**Figure 1.** The trend of Oviposition of two *T. vassilievi* populations on two CSP populations and their reciprocal crosses: (a) TT, (b) TV, (c) VT, (d) VV, (e) T×V, and (f) V×T.




**Figure 2.** Age-specific sex ratio of two *T. vassilievi* populations on two CSP populations and their reciprocal crosses: (a) TT, (b) TV, (c) VT, (d) VV, (e) T×V, and (f) V×T.

curve of *T. vassilievi* was from type I ( $H < 0.5$ ) in all treatments (Figure 3, Table 3). The life expectancy ( $e_x$ ) decreased linearly from birth to death (Figure 4). The life expectancy of *T. vassilievi* at birth and emergence, as well as under the curve area of  $e_x$  and  $N_x$  are shown in Table 3.

### Reproduction Tables for *T. vassilievi*


The reproductive parameters of *T. vassilievi* on CSP eggs are shown in Table 4. Varamin population has higher gross and net fecundity and fertility rates than Tabriz population. The crosses had higher values of these parameters than the main populations, with the highest value in T×V. The emergence rate of wasps was very high,



**Figure 3.** The survivorship curve of two *T. vassilievi* populations on two CSP populations and their reciprocal crosses: (a) TT, (b) TV, (c) VT, (d) VV, (e) T×V, and (f) V×T.

**Table 3.** Life table statistics of two *T. vassilievi* populations on two CSP populations and their reciprocal crosses under laboratory conditions.

Treatments	TT	TV	VT	VV	T×V	V×T
Entropy	0.078	0.090	0.114	0.097	0.119	0.112
Life expectancy at birth	45.9	45.1	41.8	39.55	43.75	42.05
Life expectancy at adult emergence	31.9	31.1	27.8	25.55	29.75	28.05
Under curve area of $N_x$ (Insect-day)	928	912	846	801	885	851
Under curve area of life expectancy $e_x$	1108.42	1070.47	945.75	832.91	1050.67	956.47



**Figure 4.** The life expectancy of two *T. vassilievi* populations on two CSP populations and their reciprocal crosses: (a) TT, (b) TV, (c) VT, (d) VV, (e) T×V, and (f) V×T.

ranging from 92% to 94%. The mean age of emergence was lower in Tabriz wasps and higher in Varamin than in the other treatments. The mean reproductive age for *T. vassilievi* was around 19 days. Other variables are shown in Table 4.

#### Stable Population Growth Parameters of *T. vassilievi*

Bootstrap estimates of stable population growth parameters of *T. vassilievi* on CSP eggs and their statistical comparisons are shown in Table 5. No significant difference was observed between values in columns 1

**Table 4.** Reproductive parameters of *T. vassilievi* on sunn pest eggs.

Treatments	TT	TV	VT	VV	T×V	V×T
Gross fecundity rate	201.53	212.20	229.53	232.56	262.21	250.98
Net fecundity rate	201.39	211.80	229.05	231.80	261.76	250.67
Gross fertility rate	187.20	195.60	211.89	216.52	245.29	234.19
Net fertility rate	187.09	195.26	211.48	215.85	244.97	233.96
Gross hatch rate	0.93	0.92	0.92	0.93	0.94	0.93
Mean age hatch (d)	24.67	24.23	25.97	25.28	24.70	24.95
Mean age gross fecundity (d)	19.17	19.30	19.72	19.82	19.15	19.40
Mean age net fecundity (d)	19.16	19.28	19.69	19.78	19.13	19.38
Mean age gross fertility (d)	19.04	19.11	19.53	19.61	19.01	19.22
Mean age net fertility (d)	19.03	19.09	19.50	19.57	18.99	19.21
Mean egg per day	5.17	5.58	5.89	7.05	6.24	6.44
Mean fertile egg per day	4.80	5.15	5.43	6.56	5.84	6.00
Eggs/Female/Day	6.31	6.81	8.24	9.07	8.95	5.96
Fertile eggs/Female/day	5.86	6.28	7.61	8.45	8.37	5.56

and 2, neither between columns 3 and 4. These comparisons refer to host populations and indicate the non-significant effect of the host. In contrast, comparisons between parasitoids of original populations (either between columns 1 and 3 or between columns 2 and 4) indicate significant differences in some parameters. For example, the parasitoid of Varamin population exhibited a higher level of reproduction (both GRR and  $R_0$ ) compared to Tabriz population. On the other hand, *T. vassilievi* from Varamin origin exhibited a higher rate of population increase, finite population increase, birth rate, and shorter doubling time. Crosses showed a higher reproduction rate than both original populations, with a minor advantage of V×T. In addition, their  $r_m$ -values were slightly higher than Tabriz parasitoids. Crossing between the two populations resulted in a 13.9 to 18.5% higher net reproduction rate than the maternal populations and 7.7 to 25.2% than the paternal ones. It seems that the reproductive phenotype of the progeny of outcrosses followed the maternal phenotype. The other statistics are also presented in Table 5. Estimates of stable age distribution ( $C_x$ ) showed that adults make about 1% of a stable population.

## DISCUSSION

In the present study, two geographical populations of *T. vassilievi* were compared in terms of life history parameters by considering two host populations. Outcrosses between populations with the aim of improving parasitoid fitness and breeding parasitoids with superior characteristics were also carried out. Overall, it can be stated that the parasitoids of Varamin population had a minor advantage over the Tabriz wasps. The most significant difference was in their reproductive rate, which was 8-15% higher in Varamin wasps. Increased reproduction was due to 2-days longer oviposition period in Varamin wasps. However, the daily fecundity rate was similar in both groups and did not differ significantly. Therefore, it can be concluded that wasps increase their reproduction by lengthening the oviposition period rather than increasing daily fecundity. Hence, the best way to improve the efficiency of these wasps is to select wasps with longer reproductive periods. Higher fecundity, on the other hand, reduced female longevity: Varamin females lived four days less, while the males lived one day more than the Tabriz population. The above statements were not true for crosses. The daily and total fecundity of the cross wasps increased independent of the oviposition period, as

**Table 5.** Population growth parameters of *T. vassilievi* on two populations of sunn pest and their reciprocal crosses under laboratory conditions.<sup>a</sup>

Treatments	TT	TV	VT	VV	T×V	V×T
GRR (Female/Generation)	168.21±7.24C	172.00±5.06C	188.10±5.93B	190.34±4.03B	208.38±11.06AB	216.38±4.85A
R <sub>0</sub> (Female/Generation)	168.14±7.23C	172.00±5.06C	187.95±5.96B	189.86±4.04B	208.23±11.13AB	216.22±4.90A
r <sub>m</sub> (d <sup>-1</sup> )	0.291±0.0033C	0.296±0.002BC	0.302±0.0023AB	0.303±0.0020AB	0.305±0.0032A	0.302±0.0025AB
λ (d <sup>-1</sup> )	1.337±0.0043C	1.345±0.0037BC	1.353±0.0031AB	1.354±0.0027AB	1.357±0.0044A	1.353±0.0033AB
T (d)	17.63±0.13AB	17.37±0.14BC	17.33±0.16BC	17.31±0.10C	17.48±0.13ABC	17.79±0.13A
DT (d)	2.31±0.04A	2.27±0.03AB	2.22±0.03BC	2.21±0.02BC	2.21±0.05C	2.22±0.03BC
b (d <sup>-1</sup> )	0.292±0.0033C	0.297±0.0028BC	0.303±0.0023AB	0.304±0.0020AB	0.307±0.0033A	0.303±0.0025AB
d (d <sup>-1</sup> )	0.0010±3.4×10 <sup>-5</sup> C	0.0011±3.02×10 <sup>-5</sup> BC	0.0012±2.6×10 <sup>-5</sup> AB	0.0012±2.3×10 <sup>-5</sup> A	0.0012±3.6×10 <sup>-5</sup> A	0.0012±2.8×10 <sup>-5</sup> AB

<sup>a</sup> Means bearing the same capital letters in a row are not significantly different (Bootstrap's pairwise comparisons,  $\alpha = 0.05$ ).

they exhibited 20% and 10% higher fecundity compared to Tabriz and Varamin wasps, respectively, with similar reproductive periods. This has been achieved by the introduction of new genes into the genetic pool of the original populations. The reason that males were not affected by crossing may be due to receiving only one copy of their maternal genes. Female longevity is an important qualitative indicator of parasitoids in the field. The longer life span of a parasitoid, the higher encounter to hosts, thus a wasp will have more fortune to find and exploit hosts (Suh *et al.*, 2000). However, higher reproductive effort results in the exhaustion of females themselves and shortens their lifespan (Krebs and Davies, 1993). This can be deduced by comparing the longevity of the more fecund Varamin population compared to Tabriz one. As it can be seen, Varamin wasps had higher fecundity and, at the same time, lower life span than the Tabriz wasps, which may confirm the above statement.

The average fecundity recorded for *T. vassilievi* in this study was higher than the other telenomin species. It was 98.0, 22.4, 29.6, and 63.7 for *T. biproruli*, (James, 1988), *Telenomus calvus* Johnson, (Orr *et al.*, 1986), *T. podisi* Ashmead, and *Trissolcus euschisti* (Ashmead), respectively (Yeargan, 1982). Powell and Shepard (1982) reported 88.1-141.9 broods for different isolates of *T. basalis*. The average progeny number of *T. semistriatus* was reported 88 in Turkey (Kivan and Kilic, 2006), and 210 in Varamin, Iran (Asgari, 2002). Also, it was 85 for Tabriz (Nozad Bonab, 2009), and 200 for the Varamin population of *T. grandis* (Amir-Maafi, 2000). The difference between the two populations of *T. vassilievi* was minor and not comparable to the above species. Perhaps, one reason is that the physical conditions are quite similar for both populations in the present study, while it may be deeply different in the two separate studies on a single species in the above examples. A higher number of daily fecundities may benefit augmentation programs (van Driesche and Bellows, 1996).



The sex ratio of progeny can change as parasitoid get older (Bueno *et al.*, 2008; Amir-Maafi and Parker, 2011), because, sperm reserves of the female are depleting and, as a result, insemination and, consequently, female offspring decreases (Kivan and Kiliç, 2006; Amir-Maafi and Parker, 2011). Higher female progeny benefits the scelionids, because it reduces the competition between brothers for mating with sisters (Wilson, 1961; Safavi, 1968).

Among the stable population growth parameters of *T. vassilievi*, only the gross and net reproductive rates exhibited appreciable variation among treatments. In general, cross wasps had higher reproductive rates than the original populations, which may be due to the flowing new genes in their original pool. The gross and net reproductive rates of *T. grandis* (Amir-Maafi, 2000), and *T. semistriatus* (Asgari, 2002) were 136 and 130 daughters per generation, both less than the values obtained in this study.

The intrinsic rate of population increase ( $r_m$ ) is a useful indicator of the fitness of a species or population in response to physical and/or nutritional conditions (Southwood and Henderson, 2000). This parameter can be used as a criterion for selecting natural enemies and predicting the success of bio-control agents (van Lenteren, 2003). The  $r_m$ -value of *T. vassilievi* in this study varied between 0.291 and 0.305  $d^{-1}$ , which was slightly higher in Varamin and crossed populations compared to the Tabriz population. It refers mainly to their higher fecundity. Among *Trissolcus* species, the maximum value of  $r_m$  has been reported as 0.368  $d^{-1}$  on *T. grandis* (Nozad Bonab, 2009). The  $r_m$ -value of *T. semistriatus* has been 0.226, and 0.227  $d^{-1}$  for an Iranian and a Turkish population, respectively (Asgari, 2002, Kivan and Kilic, 2006). These three species are considered as the most effective species in controlling CSP. Based on  $R_0$  and  $r_m$ -values we can rank them as *T. grandis* > *T. vassilievi* > *T. semistriatus*.

The generation Time (T) was 17.31-17.79 days for *T. vassilievi* in this study and 13.43 days for *T. grandis* (Amir-Maafi, 2000),

which can well explain why *T. grandis* is the most prevalent egg parasitoid species of CSP in Iran (Radjab and Amir Nazari, 1989). Life history data suggest the high importance of *T. vassilievi* for the biological control of CSP. A highly female-biased sex ratio, high attack rate, and longevity are positive properties for *T. vassilievi*. However, field data at different climatic conditions are essential to prove the role of *T. vassilievi* in large-scale inundation programs. Several field studies indicate the acceptable effect of some telenomins on target pests (Justo *et al.*, 1997; van Lenteren and Bueno, 2003; Asgari *et al.*, 2010; Bagheri Matin *et al.*, 2010; Asgari, 2011).

In conclusion, it can be stated that the host population had no significant effect on parasitism by *T. vassilievi*, however, fecundity was significantly higher in crosses compared to the original populations. This suggests increased fitness of progeny. Similar effects were observed on the thermal phenotypes of this species (Iranipour *et al.*, 2015). This has been demonstrated in some studies (e.g. Carson, 1968; Rasanen and Kruuk, 2007), and, nowadays, it is accepted as a scientific rule by most biologists and can be used for the artificial selection of parasitoids in the laboratory (Arakawa *et al.* 2004). The intrinsic rate of increase ( $r_m$ ) of *T. vassilievi* was less variable among treatments of this study and lied between *T. grandis* and *T. semistriatus*. The results of this study revealed that we can benefit from the intra-populations diversity of *Trissolcus* species to obtain more advantageous parasitoids via out-crossing them.

## ACKNOWLEDGEMENTS

The research facilities and funding were provided by the University of Tabriz.

## REFERENCES

1. Abdi, F., Iranipour, S. and Hejazi, M. J. 2017. Reproductive-Life Table Studies on

- Trissolcus djadetshkoe* (Hym.: Scelionidae). *Appl. Entomol. Zool.*, **85(1)**: 1-9.
2. Ahmadpour S., Iranipour, S. and Asgari, S. 2013. Effects of Superparasitism on Reproductive Fitness of *Ooencyrtus fecundus* Ferriere and Voegelé (Hym. Encyrtidae), Egg Parasitoid of Sunn Pest, *Eurygaster integriceps* Puton (Hem. Scutelleridae). *Biological Control of pests and Plant Diseases*, **2(2)**: 97-105. [in Persian]
  3. Amir-Maafi, M. 2000. An Investigation on the Host-Parasitoid System between *Trissolcus grandis* Thomson (Hym.: Scelionidae) and Sunn Pest Eggs. PhD Thesis, Faculty of Agriculture, University of Tehran, Karaj, Iran. [in Persian with English Summary].
  4. Amir-Maafi, M., Hosseini, M., Tagaddosi, M.V., Nouri, H., Haghshenas, A., Ghazi, M., Pourghz, A., Jamshidi, R., Forouzan, M. and Bagheri, S. 2010. *The Biological Control of Sunn Pest, Eurygaster integriceps* Put. (Het.: Scutelleridae) Using Egg Parasitoids. Research Project, Iranian Research Institute of Plant Protection, Tehran, Iran.
  5. Amir-Maafi, M. and Parker, B.L. 2011. Biological Parameters of the Egg Parasitoid *Trissolcus grandis* (Hym.: Scelionidae) on *Eurygaster integriceps* (Hem.: Scutelleridae). *J. Entomol. Soc. Iran.*, **30(2)**: 67-81.
  6. Arakawa, R., Miura, M. and Fujita, M. 2004. Effects of Host Species on the Body Size, Fecundity, and Longevity of *Trissolcus mitsukurii* (Hymenoptera: Scelionidae), a Solitary Egg Parasitoid of Stink Bugs. *Appl. Entomol. Zool.*, **39**: 177-181.
  7. Asgari, S. 2002. Comparative Fitness of the Eggs of *Graphosoma lineatum* (L.) (Pentatomidae) and *Eurygaster integriceps* Put. (Scutelleridae) to the Egg Parasitoid *Trissolcus semistriatus* Nees (Scelionidae). Ph.D. Thesis, Tarbiat Modares University, Tehran, Iran. [in Persian with English summary].
  8. Asgari, S. 2011. Inundative Release of the Sunn Pest Egg Parasitoid and Evaluation of Its Performance. *Proceedings of the Biological Control Development Congress in Iran*, 27-28 July, Tehran, Iran, PP. 423-428. [in Persian with English Summary].
  9. Asgari S. and Kharrazi Pakdel, A. 1998. Evaluation of Some Biological Parameters Affecting Sunn Pest Egg Parasitoid, *Trissolcus grandis* (Thom.) (Hym., Scelionidae). *Proceedings of the 13<sup>th</sup> Iranian Plant Protection Congress*, 23-27 August, Karaj, Iran, **1**: 28.
  10. Asgari, S., Safari, M. and Hassani, A.A. 2010. Mass Rearing, Releasing, and Performance of Sunn Pest Egg Parasitoids. *Proceedings of the 19<sup>th</sup> Iranian Plant Protection Congress*, 31 July-3 August, Iranian Research Institute of Plant Protection, Tehran, Iran, 72 PP.
  11. Awan, M.S., Wilson, L.T. and Hoffmann, M. P. 1990. Comparative Biology of Three Geographic Populations of *Trissolcus basalis* (Hymenoptera: Scelionidae). *Environ. Entomol.*, **19(2)**: 387-392.
  12. Bagheri Matin, S., Safavi, A., Babakfard, A and Maleki, N. 2010. Study on Efficacy of Releasing *Trissolcus grandis* Thom. (Hym: Scelionidae) on Controlling *Eurygaster integriceps* Put. (Het.: Scutelleridae) in Wheat Fields of Kermanshah Province. *Proceedings of the 19<sup>th</sup> Iranian Plant Protection Congress*, 31 July-3 August, Iranian Research Institute of Plant Protection, Tehran, Iran, 32 PP.
  13. Bazavar, A., Iranipour, S. and Karimzadeh, R. 2015. Effect of Host Unavailability Durations on Parasitism Behavior of *Trissolcus Grandis* (Hymenoptera: Scelionidae) Egg Parasitoid of Sunn Pest. *J. Appl. Res. Plant Prot.*, **4(1)**: 41-56.
  14. Benamolaei, P., Iranipour, S. and Asgari, S. 2015a. Effect of the Host Embryogenesis on Efficiency of *Trissolcus vassilievi*. *BioControl in Plant Protection*, **3(1)**: 83-100. [in Persian with English Summary]
  15. Bena molaei, P., Iranipour, S., and Asgari, S. 2015b. Biostatistics of *Trissolcus vassilievi* (Hym., Scelionidae) developed on sunn pest eggs cold-stored for different durations. *Mun. Ent. Zool.*, **10(1)**: 259-271.
  16. Bueno, R. C. O., Carneiro, T. R., Pratisoli, D., Bueno, A. D. F. and Fernandes, A. 2008. Biology and Thermal Requirements of *Telenomus remus* Reared on Fall Armyworm *Spodoptera frugiperda* Eggs. *Cienc. Rural*, **38(1)**: 1-6.
  17. Carey, J. R. 1993. Applied Demography for Biologists with Special Emphasis on Insects. Oxford University Press, UK.





18. Carson, H. L. 1968. The Population Flush and Its Genetic Consequences. In: "Population Biology and Evolution", (Ed.): Lewontin, R. C. Syracuse University Press, Syracuse, New York, PP. 123–137.
19. Iranipour, S. 2018. A Microsoft Excel Program for Bootstrap Estimates of Reproductive-Life Table Parameters. *J. Crop Prot.*, **7(3)**: 247-258.
20. Iranipour, S., Benamolaei, P., Asgari, Sh. and Michaud, J. P. 2015. Reciprocal Crosses between Two Populations of *Trissolcus vassilievi* (Mayr) (Hymenoptera: Scelionidae) Reveal Maternal Effects on Thermal Phenotypes. *Bull. Entomol. Res.*, **105**: 355-363.
21. James, D. G. 1988. Fecundity, Longevity and Overwintering of *Trissolcus biproruli* Girault (Hymenoptera: Scelionidae) a Parasitoid of *Biprorulus bibax* Breddin (Hemiptera: Pentatomidae). *J. Austral. Ent. Soc.*, **27**: 297–301.
22. Justo, H. D., Shepard B. M. and Elsey, K. D. 1997. Dispersal of the Egg Parasitoid *Trissolcus basalis* (Hymenoptera: Scelionidae) in Tomato. *J. Agric. Entomol.*, **14**: 139–149.
23. Kivan, M. and Kiliç, N. 2006. Age-Specific Fecundity and Life Table of *Trissolcus semistriatus*, an Egg Parasitoid of the Sunn Pest *Eurygaster integriceps*. *Entomol. Sci.*, **9**: 39-46.
24. Kozlov, M. A. and Kononova, S. V. 1983. Telenominae of the Fauna of the USSR (Hymenoptera, Scelionidae, Telenominae). *Leningrad Nauka Publisher*, No. 136, PP. 137–138. [in Russian]
25. Krebs, J. R. and Davies, N. B. 1993. *An Introduction to Behavioural Ecology*. Blackwell Scientific Publications.
26. Laumann, R. A., Moraes, M. C. B., Pareja, M., Alarca, G. C., Botelho, A. C., Maia, A. H. N., Leonardecz, E. and Borges, M. 2008. Comparative Biology and Functional Response of *Trissolcus* spp. Hymenoptera: Scelionidae) and Implications for Stink Bugs (Hemiptera: Pentatomidae) Biological Control. *Biol. Control.*, **44**: 32-41.
27. Lotka, A. J. 1907a. Relation between Birth Rates and Death Rates. *Science.*, **26**: 21-22.
28. Lotka, A. J. 1907b. Studies on the Mode of Growth of Material Aggregates. *Am. J. Sci.*, **4(24)**: 199-216.
29. Mele, A., Avanigadda, D. S., Ceccato, E., Olawuyi, G. B., Simoni, F., Duso, C., Scaccini, D. and Pozzebon, A. 2024. Comparative Life Tables of *Trissolcus japonicus* and *Trissolcus mitsukurii*, Egg Parasitoids of *Halyomorpha halys*. *Biol. Control.*, **195**: 105548.
30. Meyer, J. S., Ingersoll, C. G., Mac Donald, L. L. and Boyce, M. S. 1986. Estimating Uncertainty in Population Growth rates: Jackknife vs. Bootstrap Techniques. *Ecology*, **67**: 1156-1166.
31. Nouri, H., Amir-Maafi, M. and Forouzan, M. 2011. Introduction of Sunn Pest Egg Parasitoids in Qazvin, Iran. *Proceedings of the Biological Control Development Congress in Iran*, 27-28 July, Tehran, Iran, pp. 417–422. [In Persian with English Summary].
32. Nozad Bonab, Z. 2009. Effect of Temperature on Development, Fecundity and Longevity of *Trissolcus grandis* Thomson (Hym.: Scelionidae). M.Sc. Thesis, Faculty of Agriculture, University of Tabriz, Tabriz, Iran. [In Persian with English Summary]
33. Nozad Bonab, Z., Iranipour, S. and Farshbaf Pourabad, R. 2014. Demographic Parameters of Two Populations of *Trissolcus grandis* (Thomson) (Hymenoptera: Scelionidae) at Five Constant Temperatures. *J. Agric. Sci. Technol.*, **16(5)**: 969-979.
34. Orr, D. B., Russin, J. S. and Borthel, D. J. 1986. Reproductive Biology and Behavior of *Telenomus calvus* (Hymenoptera: Scelionidae), a Phoretic Egg Parasitoid of *Podisus maculiventris* (Hemiptera: Pentatomidae). *Can. Entomol.*, **118**: 1063-1072.
35. Portilla, M., Morales-Ramos, J. A., Rojas, M. G. and Blanco, C. A. 2014. Life Tables as Tools of Evaluation and Quality Control for Arthropod Mass Production. In: "Mass Production of Beneficial Organisms: Invertebrates and Entomopathogens". Academic Press, Cambridge, MA, USA, PP. 241–275.
36. Powell, J. E. and Shepard, M. 1982. Biology of Australian and United States Strains of *Trissolcus basalis*, a Parasitoid of the Green Vegetable Bug, *Nezara viridula*. *Aust. J. Ecol.*, **7**: 181-186.
37. Radjabi, G. and Amir Nazari, M. 1989. Egg Parasites of Sunn Pest in the Central Part of the Iranian Plateau. *Appl. Entomol.*

- Phytopathol.*, **56**: 1–12. [In Persian with English Summary].
38. Rasanen, K. and Kruuk, L.E.B. 2007. Maternal Effects and Evolution at Ecological Time-Scales. *Funct. Ecol.*, **21**: 408–421.
  39. Safavi, M. 1968. Etude Biologique et Ecologique des Hymenopteres Parasites des Oeufs des Punaises des Cereales. *Entomophaga.*, **13**: 381–495.
  40. Safavi, M. 1973. Etude Bio-Ecologique des Hymenoptères Parasites des Oeufs des Punaises des Cereales en Iran. Ministry of Agriculture and Natural Resources, Tehran, Iran. [in Persian].
  41. Southwood, T. R. E. and Henderson, P. A. 2000. *Ecological Methods*. 3<sup>rd</sup> Edition. Blackwell Science.
  42. Suh, C. B. C., Orr, D. B. and van Duyn, J. W. 2000. *Trichogramma* Releases in North Carolina Cotton: Why Releases Fail to Suppress Heliothine Pests. *J. Econ. Entomol.*, **93**: 1137–1145.
  43. Taghadosi, M. V., Kharrazi Pakdel, A. and Esmaili, M. M. 1993. A Comparative Study on Reproductive Potential of Different Populations of *Trissolcus grandis* Thomson (Hym., Scelionidae), on Eggs of Sunn Pest *Eurygaster integriceps* Put. (Het., Scutelleridae). In *Proceedings of the 11<sup>th</sup> Iranian Plant Protection Congress*, 27 August - 1 September, University of Guilan, Rasht, Iran, 7 PP.
  44. Teimouri, N., Iranipour, S. and Benamolaei, P. 2019. Effect of Light Intensity and Photoperiod on Development, Fecundity, and Longevity of *Trissolcus grandis* (Hym.: Platygasteridae), Egg Parasitoid of Sunn Pest, *Eurygaster integriceps* Puton (Hem.: Scutelleridae). *J. Appl. Res. Plant Prot.*, **8(3)**: 77–93.
  45. van Driesche, R. G. and Bellows Jr, T. S. 1996. *Biological Control*. Chapman and Hall, New York.
  46. van Lenteren, J. C. 2003. Quality Control and Production of Biological Control Agents: Theory and Testing Procedures. CABI Publication.
  47. van Lenteren, J.C. and Bueno, V.H.P. 2003. Augmentative Biological Control of Arthropods in Latin America. *BioControl.*, **48**: 123–139.
  48. Wilson, F. 1961. Adult Behavior in *Asolcus basalis* (Hymenoptera: Scelionidae). *Aust. J. Zool.*, **9(5)**: 737–751.
  49. Yeargan, K. V. 1982. Reproductive Capability and Longevity of the Parasitic Wasps *Telenomus podisi* and *Trissolcus euschisti*. *Ann. Entomol. Soc. Am.*, **75**: 181–183.

**افزایش شایستگی با تلاقی بین دو جمعیت زنبور (*Trissolcus vassilievi* (Hymenoptera: Scelionidae))**

شهزاد ایرانی پور، پریسا بنامولایی، و شهریار عسگری

**چکیده**

*Trissolcus vassilievi* (Hymenoptera: Scelionidae) یکی از مهم‌ترین زنبورهای پارازیتوید تخم *Eurygaster integriceps* در ایران است. در این تحقیق شایستگی دو جمعیت زنبور *T. vassilievi* روی دو جمعیت میزبان در قالب جدول‌های زندگی زادآوری مورد مطالعه قرار گرفت. دو جمعیت *T. vassilievi* یکی تبریز



(با منشاء معتدله) و دیگری ورامین (با منشاء نیمه گرمسیری) هم برای زنبور و هم برای میزبان آن لحاظ گردید. به علاوه با توجه به این که انجام تلاقی های غیرخویشاوندی بین جمعیت ها می تواند منجر به ایجاد جمعیت هایی با خصوصیات برتر شود تلاقی بین دو جمعیت روی یک میزبان واحد بررسی شد. تلاقی بین دو جمعیت موجب بروز ۱۳.۹ تا ۱۸.۵% زادآوری خالص بیشتر نسبت به جمعیت های مادری شد که به نظر می رسد بیشتر تابع فنوتیپ مادری باشد. نرخ ذاتی افزایش جمعیت اختلاف جزئی نشان داد و بین  $0.291 \pm 0.003$  تا  $0.305 \pm 0.003$  در تیمارهای مختلف متغیر بود. برتری جزئی زنبورهای ورامین نسبت به تبریز و تلاقی ها نسبت به جمعیت های اصلی مشاهده شد. این تفاوت ها می توانند در پرورش انبوه این زنبورها برای به دست آوردن پارازیتوئیدهایی با خصوصیات برتر مورد توجه قرار گیرند.